

Appropriate Testing for Pharyngitis (CWP)

New for 2023

Updated

• Members who died during the measurement year is now a required exclusion



Definition

Percentage of episodes for members age 3 years and older where the member was diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus (strep) test within 3 days prior to or 3 days after the diagnosis day (7 days total).

A higher rate indicates appropriate testing and treatment.

Plan(s) Affected	Quality Program(s) Affected	Collection and Reporting Method
Commercial	CMS Quality Rating System	Administrative
 Exchange/Marketplace 	NCQA Health Plan Ratings	Claim/Encounter Data
Medicaid		Pharmacy Data
Medicare		

Codes

The following codes can be used to close HEDIS® numerator gaps in care; they are not intended to be a directive of your billing practice.

Group A Strep Test	
CPT®/CPT II	87070, 87071, 87081, 87430, 87650, 87651, 87652, 87880
LOINC	11268-0, 17656-0, 17898-8, 18481-2, 31971-5, 49610-9, 5036-9, 60489-2, 626-2, 6557-3, 6558-1, 6559-9, 68954-7, 78012-2
SNOMED	122121004, 122205003, 122303007

Pharyngitis		
ICD-10 Diagnosis	J02.0, J02.8, J02.9, J03.00, J03.01, J03.80, J03.81, J03.90, J03.91	
SNOMED	140004, 652005,1532007, 2365002, 10351008, 11461005, 14465002, 17741008, 27878001, 31309002, 39271004, 40766000, 41582007, 43878008, 51209006, 55355000, 58031004, 59471009, 63866002, 72430001, 76651006, 78430008, 78911000, 82228008, 87326000, 90176007, 90979004, 95885008, 111816002, 126664009, 126665005, 164256007, 164260005, 186659004, 186963008, 195655000, 195656004, 195657008, 195658003, 195659006, 195660001, 195662009, 195663004, 195666007, 195667003, 195668008, 195669000, 195670004, 195671000, 195672007, 195673002, 195676005, 195677001, 195709006, 195779005, 195780008, 195782000, 195803003, 195804009, 195924009, 232399005, 232400003, 232401004, 232402006, 232403001, 232405008, 232406009, 232417005, 240444009, 240547000, 302911003, 312422001, 363746003, 405737000, 415724006, 703468005, 721586007, 878818001, 133171000119105, 10629231000119109, 10629271000119107	



Appropriate Testing for Pharyngitis (CWP)

Medications

The following antibiotic medications, in conjunction with a strep test, will meet compliance for this measure:

Drug Category	Medications	
Aminopenicillins	Amoxicillin	Ampicillin
Beta-lactamase inhibitors	Amoxicillin-clavulanate	
First generation cephalosporins	Cefadroxil Cefazolin	Cephalexin
Folate antagonist	Trimethoprim	
Lincomycin derivatives	Clindamycin	
Macrolides	AzithromycinClarithromycinErythromycin	
Natural penicillins	Penicillin G potassium Penicillin G sodium	Penicillin V potassiumPenicillin G benzathine
Quinolones	Ciprofloxacin Levofloxacin	 Moxifloxacin Ofloxacin
Second generation cephalosporins	Cefaclor Cefprozil	Cefuroxime
Sulfonamides	Sulfamethoxazole-trimethoprim	
Tetracyclines	Doxycycline Minocycline	Tetracycline
Third generation cephalosporins	Cefdinir Gefixime	CefpodoximeCeftriaxone

(Medications continued)



Appropriate Testing for Pharyngitis (CWP)

Required Exclusion(s)

Exclusion	Timeframe
Members in hospice or using hospice servicesMembers who died	- Any time during the measurement year
• HIV	- 12 months prior to or on the episode date
Malignant Neoplasms	
Malignant Neoplasms of the Skin	
Emphysema	
• COPD	
Disorders of the Immune System	



Important Notes

This measure addresses appropriate diagnosis and treatment for pharyngitis with a strep test being completed three days before or three days after the primary diagnosis and prescribed antibiotics.

A pharyngitis diagnosis can be from an outpatient, telephone, e-visit, virtual check-in, observation or emergency department visit between July 1 of the year prior to the measurement year and June 30 of the measurement year.

Medical Record Detail Including, But Not Limited to

- History and physical
- · Lab reports
- Progress notes

Tips and Best Practices to Help Close This Care Opportunity

- Please check your Patient Care Opportunity Report (PCOR) often to see members with open care opportunities. If you have questions, your UnitedHealthcare representative can help.
- Do not prescribe antibiotics until results of Group A Strep test are received.
- Always bill using the LOINC codes previously listed with your strep test submission – not local codes.
- Always use a point of care rapid Group A strep test or throat culture, when appropriate, to confirm diagnosis of pharyngitis before prescribing an antibiotic.
- Lab results can be accepted as supplemental data.
 Please contact your UnitedHealthcare representative to discuss clinical data exchange opportunities.