

Blood Pressure Control for Patients With Diabetes (BPD)

New for 2023

Added

- A direct reference code, Z51.5, for an encounter for palliative care
- Frailty exclusion now requires 2 different dates of service during the measurement year

Updated

- Members who died during the measurement year is now a required exclusion



Yes!

Supplemental Data Accepted

Definition

Percentage of members ages 18–75 with diabetes (Types 1 and 2) who have a blood pressure (BP) reading of <140/90 mmHg in the measurement year.

Plan(s) Affected	Quality Program(s) Affected	Collection and Reporting Method
<ul style="list-style-type: none"> • Commercial • Medicaid • Medicare 	<ul style="list-style-type: none"> • CMS Star Ratings • NCQA Accreditation • NCQA Health Plan Ratings 	Hybrid <ul style="list-style-type: none"> • Claim/Encounter Data • Medical Record Documentation

Codes

The following codes can be used to close HEDIS® numerator gaps in care; they are not intended to be a directive of your billing practice.

Diastolic Blood Pressure Levels

CPT®/CPT II | 3078F, 3079F, 3080*

Systolic Blood Pressure Levels

CPT®/CPT II | 3074F, 3075F, 3077*

*Please continue to code using CPT II codes for a blood pressure reading including a diastolic >90 and systolic >140, as it is important for tracking and addressing quality of care and health outcomes.

HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA). CPT® is a registered trademark of the American Medical Association. UnitedHealthcare will make the final determination regarding reimbursement upon receipt of a claim. Submitting a claim with a code included in this document is not a guarantee of payment. Payment of covered services is contingent upon coverage within an individual member's benefit plan, your eligibility for payment, any claim processing requirements, and your participation agreement with UnitedHealthcare.

Blood Pressure Control for Patients With Diabetes (BPD)

Required Exclusion(s)

Exclusion	Timeframe
<ul style="list-style-type: none"> Members in hospice or using hospice services Members receiving palliative care Members who died 	Any time during the measurement year
Members who have no diagnosis of diabetes and have a diagnosis of polycystic ovarian syndrome, gestational diabetes or steroid-induced diabetes	During the measurement year or year prior
<p>Members ages 66 and older as of December 31 of the measurement year who had 2 diagnoses of frailty on different dates of service and advanced illness.* Advanced illness is indicated by one of the following:</p> <ul style="list-style-type: none"> Two or more outpatient, observation, emergency room, telephone, e-visits, virtual check-ins or non-acute inpatient encounters or discharge(s) on separate dates of service with a diagnosis of advanced illness One or more acute inpatient encounter(s) with a diagnosis of advanced illness One or more acute inpatient discharge(s) with a diagnosis of advanced illness on the discharge claim Dispensed a dementia medication: Donepezil, Donepezil-memantine, galantamine, rivastigmine or memantine 	<p>Frailty diagnoses must be in the measurement year on 2 different dates of service</p> <p>Advanced illness diagnosis must be in the measurement year or year prior to the measurement year</p>
<p>Medicare members ages 66 and older as of December 31 of the measurement year who are either:</p> <ul style="list-style-type: none"> Enrolled in an Institutional Special Needs Plan (I-SNP) Living long term in an institution* 	Any time during the measurement year

*Supplemental and medical record data may not be used for the frailty with advanced illness or institutional living exclusions.

Blood Pressure Control for Patients With Diabetes (BPD)



Important Notes

	Test, Service or Procedure to Close Care Opportunity	Medical Record Detail Including, But Not Limited To
<ul style="list-style-type: none"> • BP reading must be performed within the measurement year — last BP result of the year is the one measured. • BP readings taken on the same day the member receives a common low-intensity or preventive procedure can be used. Examples include, but aren't limited to: <ul style="list-style-type: none"> – Eye exam with dilating agents – Injections (e.g., allergy, Depo-Provera®, insulin, lidocaine, steroid, testosterone toradol or vitamin B-12) – Intrauterine device (IUD) insertion – Tuberculosis (TB) test – Vaccinations – Wart or mole removal 	<p>BP reading taken or reported and recorded during the measurement year via outpatient visits, telephone or telehealth visits, e-visits, virtual check-ins, or non-acute inpatient visits. Member-reported BP readings must be taken using a digital device in any of these visit settings and documented in member's medical record (must note "digital device").</p>	<ul style="list-style-type: none"> • Consultation reports • Diabetic flow sheets • Progress notes • Vitals sheet

(Important Notes continued)

Blood Pressure Control for Patients With Diabetes (BPD)



Important Notes

	Test, Service or Procedure to Close Care Opportunity	Medical Record Detail Including, But Not Limited To
<ul style="list-style-type: none"> • BP readings taken in the following situations will not count toward compliance: <ul style="list-style-type: none"> - During an acute inpatient stay or an emergency department visit - On the same day as a diagnostic test, or diagnostic or therapeutic procedure that requires a change in diet or medication on or one day before the day of the test or procedure – with the exception of a fasting blood test. Examples include, but are not limited to: <ul style="list-style-type: none"> • Colonoscopy • Dialysis, infusions and chemotherapy • Nebulizer treatment with albuterol • BP readings taken by a member using a non-digital device, e.g., manual blood pressure cuff and stethoscope, do not meet numerator compliance. 		<ul style="list-style-type: none"> • Consultation reports • Diabetic flow sheets • Progress notes • Vitals sheet

Blood Pressure Control for Patients With Diabetes (BPD)

Tips and Best Practices to Help Close This Care Opportunity

- **Please check your Patient Care Opportunity Report (PCOR) often to see members with open care opportunities.** If you have questions, your UnitedHealthcare representative can help.
- It is important to document patient reported vitals in the official medical record when conducting telehealth, telephone or online assessment visits. Please encourage patients to use a digital device to track and report their BP during every visit.
- **Always list the date of service and BP reading together.**
 - If BP is listed on the vital flow sheet, it must have a date of service.
- Members who have an elevated BP during an office visit in Aug., Sept. or Oct. should be brought back in for a follow-up visit before Dec. 31.
- Talk with members about what a lower goal is for a healthy BP reading.
 - For example: 130/80 mmHg
- Remind members who are NPO for a fasting lab they should continue to take their anti-hypertensive medications with a sip of water on the morning of their appointment.
- If your office uses manual blood pressure cuffs, don't round up the BP reading.
 - For example: 138/89 mmHg rounded to 140/90 mmHg
- If a member's initial BP reading is elevated at the start of a visit, you can take multiple readings during the same visit and use the lowest diastolic and lowest systolic to document the overall reading. Retake the member's BP after they've had time to rest.
 - For example: **If a member's first BP reading was 160/80 mmHg and the second reading was 120/90 mmHg, use the 120 systolic of the second reading and the 80 diastolic of the first reading to show a BP result of 120/80 mmHg.**
- If your office submits CCDs to UnitedHealthcare via our clinical data exchange program, please ensure the CCD function within your EMR system is set up to send CPT II Codes in the extract.
- The use of CPT® Category II codes helps UnitedHealthcare identify clinical outcomes such as diastolic and systolic readings. It can also reduce the need for some chart review.
- BP readings can be accepted as supplemental data, reducing the need for some chart review. Please contact your UnitedHealthcare representative to discuss clinical data exchange opportunities.
- Sharing member demographic data is critical to understanding the cultural, linguistic and social needs of those we serve and decreasing health inequities across the care continuum. This data can include, but is not limited to, race, ethnicity, language, sexual orientation, gender identity, pronouns, sex assigned at birth and disability status. As part of UnitedHealthcare's clinical structured data exchange program, we encourage you to include this demographic data with any structured data file or CCD. Your information is confidential. UnitedHealthcare will keep personally identifiable information confidential and won't disclose any information without your written consent.